



Smith, M. D., Ben-Shlomo, Y., & Henderson, E. (2019). How often are patients with progressive supranuclear palsy really falling? *Journal of Neurology*, 266(8), 2073-2074. <https://doi.org/10.1007/s00415-019-09375-0>

Peer reviewed version

Link to published version (if available):
[10.1007/s00415-019-09375-0](https://doi.org/10.1007/s00415-019-09375-0)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Springer Nature at <https://link.springer.com/article/10.1007%2Fs00415-019-09375-0>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

How often are patients with Progressive Supranuclear Palsy really falling?

Matthew D. Smith¹ BM BS BSc MRCP, Yoav Ben-Shlomo² MB BS PhD, Emily Henderson² MBChB MRCP PhD

1) North Bristol NHS Trust 2) School of Social and Community Medicine, University of Bristol

Corresponding author:

Matthew Smith, North Bristol NHS Trust, Bristol, BS10 5NB, matthew.smith1@doctors.org.uk

+44 7454 011069

Word Count: 529

Key Words: *PSP, Progressive supranuclear palsy, falls, gait, balance*

Progressive supranuclear palsy (PSP) is a neurodegenerative condition marked by supranuclear ophthalmoplegia, pseudobulbar palsy, and gait instability leading to repeated falls. We undertook a systematic review to determine how the rate of falls was measured in PSP studies; a cardinal feature of the condition. We searched Pubmed using following the search terms,: ("progressive supranuclear palsy" OR PSP OR "Steele-Richardson-Olszewski syndrome") AND falls), as well as performing an individual search of the MeSH terms "progressive supranuclear palsy AND accidental falls", up to March 2019. Papers were included if they reported the incidence of falling and were in English. All study designs were included (observational or interventional) as long as they included more than one participant.

Thirty papers were identified with a mean number of participants of 47.6 (range 8-350). Study designs ranged from prospective and retrospective cohorts to small scale trials.

Only two studies documented an absolute number of falls using a falls diary (the established gold standard)[1], or mean number of falls per patient. A study by Zwergal *et al.* reported a total of 195 falls occurring in 16 patients (mean 11.8 (range 1-40) over a six month period [2]. Another study reported by Clerici *et al.* demonstrated a mean of 7 and 8 falls over a one month period from 12 patients at baseline prior to allocation to two rehabilitation programmes [3]. A Further paper (Srulijes *et al.*) reported a quantitative falls rate of 46.2 per year for a whole cohort in a comparative study of PSP patients (n=12) compared to other neurodegenerative diseases. [4] A selection of other studies quantified falls prospectively but did not report quantitative data for falls values at publication. The majority of studies used either medical record review or a semi-quantitative ordinal scale as found in the PSPRS (Progressive Supranuclear Palsy Rating Scale) or UPDRS (Unified Parkinson's Disease Rating Scale) [5,6]. Patients were usually divided into arbitrary groups such as "frequent" or "infrequent" fallers, with inconsistent cut-offs. Where medical records were reviewed, a binary variable (presence or absence of falls) was usually recorded.

It is surprising that few studies have sought to provide prospective data on fall frequency given that falls are a major feature in PSP. The PSPRS captures falls as an ordinal variable (0=None in the past year, 1= <1 per month, 2=1-4 per month, 3=5-30 per month, 4 > 30 per month or chairbound). This can be used for comparing group differences but does not provide an absolute difference in falls rate and has an unbounded top category. Additionally, the score is retrospective relying on patient/carer recall with a greater degree of measurement error and potential recall bias. However, where retrospective studies are performed and falls data collected, we appreciate that collecting absolute data retrospectively may also be obscured by recall bias making a truly accurate picture hard to obtain. We advocate that future PSP studies should collect falls data using the current gold standard ProFANE (Prevention of Falls Network Europe) methods [1], and that a prospective approach is best. This will provide more useful and reliable data on falls in this high risk population and enable better designed intervention studies to be undertaken to tackle this debilitating problem.

Conflict of interest statement:

On behalf of all authors, the corresponding author states that there is no conflict of interest.

References:

- [1] Lamb SE, Jørstad-Stein EC, Hauer K, Becker C. Development of a common outcome data set for fall injury prevention trials: the Prevention of Falls Network Europe consensus. *Journal of the American Geriatrics Society*. 2005 Sep 1;53(9):1618-22.
- [2] Zwergal A, la Fougère C, Lorenzl S, Rominger A, Xiong G, Deutschenbaur L, Linn J, Krafczyk S, Dieterich M, Brandt T, Strupp M, Bartenstein P, Jahn K. Postural imbalance and falls in PSP correlate with functional pathology of the thalamus. *Neurology*. 2011 Jul 12;77(2):101-9.
- [3] Clerici I, Ferrazzoli D, Maestri R, Bossio F, Zivi I, Canesi M, Pezzoli G, Frazzitta G. Rehabilitation in progressive supranuclear palsy: Effectiveness of two multidisciplinary treatments. *PLoS One*. 2017 Feb 3;12(2):e0170927.
- [4] Srulijes K, Klenk J, Schwenk M, Schatton C, Schwickert L, Teubner-Liepert K, Meyer M, K C S, Maetzler W, Becker C, Synofzik M. Fall Risk in Relation to Individual Physical Activity Exposure in Patients with Different Neurodegenerative Diseases: a Pilot Study. *Cerebellum*. 2019 Jan 8. doi: 10.1007/s12311-018-1002-x. [Epub ahead of print]
- [5] Golbe LI, Ohman-Strickland PA. A clinical rating scale for progressive supranuclear palsy. *Brain*. 2007 Jun;130(Pt 6):1552-65.
- [6] Goetz CG, Fahn S, Martinez-Martin P, Poewe W, Sampaio C, Stebbins GT, Stern MB, Tilley BC, Dodel R, Dubois B, Holloway R, Jankovic J, Kulisevsky J, Lang AE, Lees A, Leurgans S, LeWitt PA, Nyenhuis D, Olanow CW, Rascol O, Schrag A, Teresi JA, Van Hilten JJ, LaPelle N.. Movement Disorder Society-sponsored revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): Process, format, and clinimetric testing plan. *Movement Disorders*. 2007 22 (1): 41–47.